

# A NEW APPROACH TO THE FOURIER ANALYSIS ON SEMI-DIRECT PRODUCTS OF GROUPS

ARASH GHAANI FARASHAHI

**ABSTRACT.** Let  $H$  and  $K$  be locally compact groups and also  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \ltimes_\tau K$  be the semi-direct product of  $H$  and  $K$  with respect to the continuous homomorphism  $\tau$ . This paper presents a novel approach to the Fourier analysis of  $G_\tau$ , when  $K$  is abelian. We define the  $\tau$ -dual group  $G_{\hat{\tau}}$  of  $G_\tau$  as the semi-direct product  $H \ltimes_{\hat{\tau}} \hat{K}$ , where  $\hat{\tau} : H \rightarrow \text{Aut}(\hat{K})$  defined via (3.1). We prove a Ponterjagin duality Theorem and also we study  $\tau$ -Fourier transforms on  $G_\tau$ . As a concrete application we show that how these techniques apply for the affine group and also we compute the  $\tau$ -dual group of Euclidean groups and the Weyl-Heisenberg groups.

## 1. Introduction

Theory of Fourier analysis is the basic and fundamental step to extend the approximation theory on algebraic structures. Classical Fourier analysis on  $\mathbb{R}^n$  and also it's standard extension for locally compact abelian groups play an important role in approximation theory and also time-frequency analysis. For more on this topics we refer the readers to [3] or [4]. Passing through the harmonic analysis of abelian groups to the harmonic analysis of non-abelian groups we loose many concepts of Fourier analysis on locally compact abelian groups. If we assume that  $G$  is unimodular and type I locally compact group, then still Fourier analysis on  $G$  can be used. Theory of Fourier analysis on non-abelian, unimodular and type I groups was completely studied by Lipsman in [9] and also Dixmier in [2] or [8].

Although theory of standard non-abelian Fourier analysis is a strong theory but it is not numerical computable, so it is not an appropriate tools in the view points of time-frequency analysis or physics and engineering applications. This lake persists us to have a new approach to the theory of Fourier analysis on non-abelian groups.

Many non-abelian groups which play important roles in general theory of time-frequency analysis or mathematical physics such as the affine group or Heisenberg group can be considered as a semi-direct products of some locally compact groups  $H$  and  $K$  with respect to a continuous homomorphism  $\tau : H \rightarrow \text{Aut}(K)$  in which  $K$  is abelain.

In this paper which contains 5 sections, section 2 devoted to fix notations and also a summary of harmonic analysis on locally compact groups and semi-direct product of locally compact groups  $H$  and  $K$  with respect to the continuous homomorphism  $\tau : H \rightarrow \text{Aut}(K)$ . In section 3 we assume that  $K$  is abelian and also we define the  $\tau$ -dual group  $G_{\hat{\tau}}$  of  $G_\tau = H \ltimes_\tau K$  as the semi direct products of  $H$  and  $\hat{K}$  with respect to the continuous homomorphism  $\hat{\tau} : H \rightarrow \text{Aut}(\hat{K})$ , where  $\hat{\tau}_h(\omega) := \omega \circ \tau_{h^{-1}}$ . It is also shown that the  $\hat{\tau}$ -dual group  $G_{\hat{\tau}}$  of  $G_{\hat{\tau}} = H \ltimes_{\hat{\tau}} \hat{K}$  and  $G_\tau$  are isomorphic, which can be considered as a generalization of the Ponterjagin duality Theorem.

In the sequel, in section 4 we define  $\tau$ -Fourier transform of  $f \in L^1(G_\tau)$  and we study it's basic  $L^2$ -properties such as the Plancherel theorem. We also prove an inversion formula for the  $\tau$ -Fourier transform.

As well as, finally in section 5 as examples we show that how this extension techniques can be used for various types of semi-direct products of group such as the affine group, the Euclidean groups and the Weyl-Heisenberg groups.

---

2000 *Mathematics Subject Classification.* Primary 22A38, 22D45.

*Key words and phrases.* semi-direct products of locally compact groups,  $\tau$ -dual group,  $\tau$ -Fourier transform, Parseval formula, Plancherel Theorem, inversion formula.

E-mail addresses: ghaanifarashahi@hotmail.com (Arash Ghaani Farashahi).

## 2. Preliminaries and notations

Let  $H$  and  $K$  be locally compact groups with identity elements  $e_H$  and  $e_K$  respectively and left Haar measures  $dh$  and  $dk$  respectively and also let  $\tau : H \rightarrow \text{Aut}(K)$  be a homomorphism such that the map  $(h, k) \mapsto \tau_h(k)$  from  $H \times K$  onto  $K$  be continuous, where  $\text{Aut}(K)$  is the group of all topological group automorphisms of  $K$  onto  $K$ . There is a natural topology, sometimes called Braconnier topology, turning  $\text{Aut}(K)$  into a Hausdorff topological group (not necessarily locally compact), which is defined by the sub-base of identity neighbourhoods

$$(2.1) \quad \mathcal{B}(F, U) = \{\alpha \in \text{Aut}(K) : \alpha(k), \alpha^{-1}(k) \in Uk \ \forall k \in F\},$$

where  $F \subseteq K$  is compact and  $U \subseteq K$  is an identity neighbourhood and also continuity of a homomorphism  $\tau : H \rightarrow \text{Aut}(K)$  is equivalent to the continuity of the map  $(h, k) \mapsto \tau_h(k)$  from  $H \times K$  onto  $K$  (see [7]). The semi-direct product  $G_\tau = H \ltimes_\tau K$  is a locally compact topological group with underlying set  $H \times K$  which equipped with product topology and group operation is defined by

$$(2.2) \quad (h, k) \ltimes_\tau (h', k') := (hh', k\tau_h(k')) \quad \text{and} \quad (h, k)^{-1} := (h^{-1}, \tau_{h^{-1}}(k^{-1})).$$

If  $H_1 := \{(h, e_K) : h \in H\}$  and  $K_1 := \{(e_H, k) : k \in K\}$ , then  $K_1$  is a closed normal subgroup and  $H_1$  is a closed subgroup of  $G_\tau$ . The left Haar measure of  $G_\tau$  is  $d\mu_{G_\tau}(h, k) = \delta(h)dhdk$  and also  $\Delta_{G_\tau}(h, k) = \delta(h)\Delta_H(h)\Delta_K(k)$ , where the positive and continuous homomorphism  $\delta : H \rightarrow (0, \infty)$  is given by (Theorem 15.29 of [5])

$$(2.3) \quad dk = \delta(h)d(\tau_h(k)).$$

From now on, for all  $p \geq 1$  we denote by  $L^p(G_\tau)$  the Banach space  $L^p(G_\tau, \mu_{G_\tau})$  and also  $L^p(K)$  stands for  $L^p(K, dk)$ . When  $f \in L^p(G_\tau)$ , for a.e.  $h \in H$  the function  $f_h$  defined on  $K$  via  $f_h(k) := f(h, k)$  belongs to  $L^p(K)$  (see [4]).

If  $K$  is a locally compact abelian group, due to Corollary 3.6 of [3] all irreducible representations of  $K$  are one-dimensional. Thus, if  $\pi$  be an irreducible unitary representation of  $K$  we have  $\mathcal{H}_\pi = \mathbb{C}$  and also according to the Shur's Lemma, there exists a continuous homomorphism  $\omega$  of  $K$  into the circle group  $\mathbb{T}$  such that for each  $k \in K$  and  $z \in \mathbb{C}$  we have  $\pi(k)(z) = \omega(k)z$ . Such continuous homomorphisms are called characters of  $K$  and the set of all characters of  $K$  denoted by  $\widehat{K}$ . If  $\widehat{K}$  equipped by the topology of compact convergence on  $K$  which coincides with the  $w^*$ -topology that  $\widehat{K}$  inherits as a subset of  $L^\infty(K)$ , then  $\widehat{K}$  with respect to the dot product of characters is a locally compact abelian group which is called the dual group of  $K$ . The linear map  $\mathcal{F}_K : L^1(K) \rightarrow \mathcal{C}(\widehat{K})$  defined by  $v \mapsto \mathcal{F}_K(v)$  via

$$(2.4) \quad \mathcal{F}_K(v)(\omega) = \widehat{v}(\omega) = \int_K v(k) \overline{\omega(k)} dk,$$

is called the Fourier transform on  $K$ . It is a norm-decreasing  $*$ -homomorphism from  $L^1(K)$  to  $\mathcal{C}_0(\widehat{K})$  with a uniformly dense range in  $\mathcal{C}_0(\widehat{K})$  (Proposition 4.13 of [3]). If  $\phi \in L^1(\widehat{K})$ , the function defined a.e. on  $K$  by

$$(2.5) \quad \check{\phi}(x) = \int_{\widehat{K}} \phi(\omega) \omega(x) d\omega,$$

belongs to  $L^\infty(K)$  and also for all  $f \in L^1(K)$  we have the following orthogonality relation (Parseval formula);

$$(2.6) \quad \int_K f(k) \overline{\check{\phi}(k)} dk = \int_{\widehat{K}} \widehat{f}(\omega) \overline{\phi(\omega)} d\omega.$$

The Fourier transform (2.4) on  $L^1(K) \cap L^2(K)$  is an isometric and it extends uniquely to a unitary isomorphism from  $L^2(K)$  to  $L^2(\widehat{K})$  (Theorem 4.25 of [3]) and also each  $v \in L^1(K)$  with  $\widehat{v} \in L^1(\widehat{K})$  satisfies the following Fourier inversion formula (Theorem 4.32 of [3]);

$$(2.7) \quad v(k) = \int_{\widehat{K}} \widehat{v}(\omega) \omega(k) d\omega \text{ for a.e. } k \in K.$$

### 3. $\tau$ -Dual group

We recall that for a locally compact non-abelian group  $G$ , the standard dual space  $\widehat{G}$  is defined as the set of all unitary equivalence classes of all irreducible unitary representations of  $G$ . There is a topology on  $\widehat{G}$  called Fell topology. But  $\widehat{G}$  with respect to the Fell topology is not a locally compact group in general setting (see [3]). On the other hand elements of  $\widehat{G}$  are equivalence classes of irreducible unitary representations of  $G$  and so from computational view points there are not numerical applicable. In this section we associate to any semi-direct product group  $H \ltimes_\tau K$  with  $K$  abelian, a  $\tau$ -dual structure (group) which is actually a locally compact group.

Let  $H$  be a locally compact group and  $K$  be a locally compact abelian group also let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \ltimes_\tau K$ . For all  $h \in H$  and  $\omega \in \widehat{K}$  define the action  $H \times \widehat{K} \rightarrow \widehat{K}$  via

$$(3.1) \quad \omega_h := \omega \circ \tau_{h^{-1}},$$

where  $\omega_h(k) = \omega(\tau_{h^{-1}}(k))$  for all  $k \in K$ . If  $\omega \in \widehat{K}$  and  $h \in H$  we have  $\omega_h \in \widehat{K}$ , because for all  $k, s \in K$  we have

$$\begin{aligned} \omega_h(k) &= \omega \circ \tau_{h^{-1}}(k) \\ &= \omega(\tau_{h^{-1}}(k)) \\ &= \omega(\tau_{h^{-1}}(k)\tau_{h^{-1}}(s)) \\ &= \omega(\tau_{h^{-1}}(k))\omega(\tau_{h^{-1}}(s)) = \omega_h(k)\omega_h(s). \end{aligned}$$

In the following proposition we find a suitable relation about the Plancherel measure of  $\widehat{K}$  and also the action of  $H$  on  $\widehat{K}$  due to (3.1).

**Proposition 3.1.** *Let  $K$  be an abelian group and  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism. The Plancherel measure  $d\omega$  on  $\widehat{K}$  for all  $h \in H$  satisfies*

$$(3.2) \quad d\omega_h = \delta(h)d\omega,$$

where  $\delta : H \rightarrow (0, \infty)$  is the positive continuous homomorphism given by  $dk = \delta(h)d\tau_h(k)$ .

*Proof.* Let  $h \in H$  and also  $v \in L^1(K)$ . Using (2.3) we have  $v \circ \tau_h \in L^1(K)$  with  $\|v \circ \tau_h\|_{L^1(K)} = \delta(h)\|v\|_{L^1(K)}$ , because

$$\begin{aligned} \|v \circ \tau_h\|_{L^1(K)} &= \int_K |v \circ \tau_h(k)| dk \\ &= \int_K |v(\tau_h(k))| dk \\ &= \int_K |v(k)| d\tau_{h^{-1}}(k) \\ &= \delta(h) \int_K |v(k)| dk = \delta(h)\|v\|_{L^1(K)}. \end{aligned}$$

Thus, for all  $\omega \in \widehat{K}$  we achieve

$$\begin{aligned} \widehat{v \circ \tau_h}(\omega) &= \int_K v(\tau_h(k)) \overline{\omega(k)} dk \\ &= \int_K v(k) \overline{\omega_h(k)} d(\tau_{h^{-1}}(k)) \\ &= \delta(h) \int_K v(k) \overline{\omega_h(k)} dk = \delta(h) \widehat{v}(\omega_h). \end{aligned}$$

Now let  $v \in L^1(K) \cap L^2(K)$ . According to the Plancherel theorem (Theorem 4.25 of [3]) and also preceding calculation, for all  $h \in H$  we get

$$\begin{aligned}
\int_{\widehat{K}} |\widehat{v}(\omega)|^2 d\omega_h &= \int_{\widehat{K}} |\widehat{v}(\omega_{h^{-1}})|^2 d\omega \\
&= \delta(h)^2 \int_{\widehat{K}} |\widehat{v \circ \tau_{h^{-1}}}(\omega)|^2 d\omega \\
&= \delta(h)^2 \int_K |v \circ \tau_{h^{-1}}(k)|^2 dk \\
&= \delta(h)^2 \int_K |v(k)|^2 d(\tau_h(k)) \\
&= \delta(h) \int_K |v(k)|^2 dk = \int_{\widehat{K}} |\widehat{v}(\omega)|^2 \delta(h) d\omega,
\end{aligned}$$

which implies (3.2). □

Now using the action defined in (3.1) we define  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{K})$  via  $h \mapsto \widehat{\tau}_h$ , where

$$(3.3) \quad \widehat{\tau}_h(\omega) := \omega_h = \omega \circ \tau_{h^{-1}}.$$

According to (3.3) for all  $h \in H$  we have  $\widehat{\tau}_h \in \text{Aut}(\widehat{K})$ . Because, if  $k \in K$  and  $h \in H$  then for all  $\omega, \eta \in \widehat{K}$  we have

$$\begin{aligned}
\widehat{\tau}_h(\omega.\eta)(k) &= (\omega.\eta)_h(k) \\
&= (\omega.\eta) \circ \tau_{h^{-1}}(k) \\
&= \omega.\eta(\tau_{h^{-1}}(k)) \\
&= \omega(\tau_{h^{-1}}(k))\eta(\tau_{h^{-1}}(k)) \\
&= \omega_h(k)\eta_h(k) = \widehat{\tau}_h(\omega)(k)\widehat{\tau}_h(\eta)(k).
\end{aligned}$$

Also  $h \mapsto \widehat{\tau}_h$  is a homomorphism from  $H$  into  $\text{Aut}(\widehat{K})$ , cause if  $h, t \in H$  then for all  $\omega \in \widehat{K}$  and also  $k \in K$  we have

$$\begin{aligned}
\widehat{\tau}_{th}(\omega)(k) &= \omega_{th}(k) \\
&= \omega(\tau_{(th)^{-1}}(k)) \\
&= \omega(\tau_{h^{-1}}\tau_{t^{-1}}(k)) \\
&= \omega_h(\tau_{t^{-1}}(k)) \\
&= \widehat{\tau}_h(\omega)(\tau_{t^{-1}}(k)) = \widehat{\tau}_t[\widehat{\tau}_h(\omega)](k).
\end{aligned}$$

Thus, via an algebraic viewpoint we can consider the semi-direct product of  $H$  and  $\widehat{K}$  with respect to the homomorphism  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{K})$ . Due to (2.2),  $\widehat{\tau}$ -dual group operation for all  $(h, \omega), (t, \eta) \in G_{\widehat{\tau}} = H \ltimes_{\widehat{\tau}} \widehat{K}$  is

$$(3.4) \quad (h, \omega) \ltimes_{\widehat{\tau}} (t, \eta) = (ht, \omega.\eta_h).$$

Now we are in the position to prove the following fundamental theorem.

**Theorem 3.2.** *Let  $H$  and  $K$  be locally compact groups with  $K$  abelian,  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and also let  $\delta : H \rightarrow (0, \infty)$  be the positive continuous homomorphism given via  $dk = \delta(h)d\tau_h(k)$ . The homomorphism  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{K})$  defined in (3.1) is continuous and so that the semi-direct product  $H \ltimes_{\widehat{\tau}} \widehat{K}$  is a locally compact group with the left Haar measure  $d\mu_{G_{\widehat{\tau}}}(h, \omega) = \delta(h)^{-1}dh d\omega$ .*

*Proof.* For  $\alpha \in \text{Aut}(K)$  let  $\widehat{\alpha} \in \text{Aut}(\widehat{K})$  be given for all  $\omega \in \widehat{K}$  by  $\widehat{\alpha}(\omega) := \omega \circ \alpha^{-1}$  where for all  $k \in K$  we have  $\omega \circ \alpha^{-1}(k) = \omega(\alpha^{-1}(k))$ . Due to Theorem 26.9 and also Theorem 26.5 of [5] the mapping  $\widehat{\cdot} : \text{Aut}(K) \rightarrow \text{Aut}(\widehat{K})$  defined by  $\alpha \mapsto \widehat{\alpha}$  is a topological group isomorphism and so it is continuous. According to the following diagram

$$(3.5) \quad H \xrightarrow{\tau} \text{Aut}(K) \xrightarrow{\widehat{\cdot}} \text{Aut}(\widehat{K}),$$

the homomorphism  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{K})$  defined in (3.1) is continuous. Thus, the semi-direct product  $H \ltimes_{\widehat{\tau}} \widehat{K}$  is a locally compact group and also Proposition 3.1 shows that  $\delta(h)^{-1}dh d\omega$  is a left Haar measure for  $H \ltimes_{\widehat{\tau}} \widehat{K}$ .  $\square$

The semi-direct product  $G_{\widehat{\tau}} = H \ltimes_{\widehat{\tau}} \widehat{K}$  mentioned in Theorem 3.2, called the  $\tau$ -dual group of  $G_{\tau} = H \ltimes_{\tau} K$ . The most important advantage of this definition as a kind of a dual space for semi direct product of locally compact groups is that its elements are numerical computable and also this dual space is merely a locally compact group. It is worthwhile to note that, when  $H$  is the identity group, the  $\tau$ -dual group of  $G_{\tau} = K$  coincides with the usual dual group  $\widehat{K}$  of  $K$ . When  $K$  is abelian locally compact group and  $\tau : H \rightarrow \text{Aut}(K)$  is a continuous homomorphism, we call  $K$  as the Fourier factor of the semi-direct product  $G_{\tau} = H \ltimes_{\tau} K$ .

Due to the Pontrjagin duality theorem (Theorem 4.31 of [3]), each  $k \in K$  defines a character  $\widehat{k}$  on  $\widehat{K}$  via  $\widehat{k}(\omega) = \omega(k)$  and also the map  $k \mapsto \widehat{k}$  is a topological group isomorphism from  $K$  onto  $\widehat{\widehat{K}}$ . Via the same method as introduced in (3.1) the  $\widehat{\tau}$ -dual group operation, for all  $(h, \widehat{k})$  and  $(t, \widehat{s})$  in  $G_{\widehat{\tau}} = H \ltimes_{\widehat{\tau}} \widehat{\widehat{K}}$  is

$$(3.6) \quad (h, \widehat{k}) \ltimes_{\widehat{\tau}} (t, \widehat{s}) = (ht, \widehat{k\widehat{\tau}_h(\widehat{s})}),$$

where  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{\widehat{K}})$  is given by

$$(3.7) \quad \widehat{\tau}_h(\widehat{k})(\omega) = \omega_{h^{-1}}(k),$$

for all  $\omega \in \widehat{\widehat{K}}$  and  $(h, k) \in G_{\tau}$ . Because, due to (3.3) we have

$$\begin{aligned} \widehat{\tau}_h(\widehat{k})(\omega) &= \widehat{k} \circ \widehat{\tau}_{h^{-1}}(\omega) \\ &= \widehat{k}(\widehat{\tau}_{h^{-1}}(\omega)) \\ &= \widehat{k}(\omega_{h^{-1}}) = \omega_{h^{-1}}(k). \end{aligned}$$

In the sequel we prove a type of Pontrjagin duality theorem for  $\tau$ -dual group of semi direct product of groups. But first we prove a short lemma.

**Lemma 3.3.** *Let  $K$  be an abelian group and  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism also let  $G_{\tau} = H \ltimes_{\tau} K$ . Then, for all  $(h, k) \in G_{\tau}$  we have*

$$(3.8) \quad \widehat{\tau_h(k)} = \widehat{\tau}_h(\widehat{k}).$$

*Proof.* Let  $(h, k) \in G_{\tau}$  and also let  $\omega \in \widehat{\widehat{K}}$ . Using duality notation and also (3.7) we have

$$\begin{aligned} \widehat{\tau_h(k)}(\omega) &= \omega(\tau_h(k)) \\ &= \omega \circ \tau_h(k) \\ &= \omega_{h^{-1}}(k) = \widehat{\tau}_h(\widehat{k})(\omega). \end{aligned}$$

$\square$

Next theorem gives us a subtle topological group isomorphism form  $G_{\tau}$  onto  $G_{\widehat{\tau}}$ . In fact, the next theorem can be considered as the Pontrjagin duality theorem for  $\tau$ -dual group of semi-direct product of groups.

**Theorem 3.4.** *Let  $H$  be a locally compact group,  $K$  be a locally compact abelian group and  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism also let  $G_{\tau} = H \ltimes_{\tau} K$ . The map  $\Theta : G_{\tau} \rightarrow G_{\widehat{\tau}}$  defined by*

$$(3.9) \quad (h, k) \mapsto \Theta(h, k) := (h, \widehat{k}),$$

*is a topological group isomorphism.*

*Proof.* First we show that  $\Theta$  is a homomorphism. Let  $(h, k), (t, s)$  in  $G_\tau$ . Since the map  $k \mapsto \widehat{k}$  is a homomorphism and also using Lemma 3.3 we have

$$\begin{aligned}\Theta((h, k) \times_\tau (t, s)) &= \Theta(ht, k\tau_h(s)) \\ &= \left(ht, \widehat{k\tau_h(s)}\right) \\ &= \left(ht, \widehat{\widehat{k}\tau_h(s)}\right) \\ &= \left(ht, \widehat{\widehat{k}\tau_h}(\widehat{s})\right) \\ &= (h, \widehat{k}) \times_{\widehat{\tau}} (t, \widehat{s}) = \Theta(h, k) \times_{\widehat{\tau}} \Theta(t, s).\end{aligned}$$

Now using Pontrjagin Theorem (Theorem 4.31 of [3]), the map  $k \mapsto \widehat{k}$  is a topological group isomorphism from  $K$  onto  $\widehat{K}$  which implies that the map  $\Theta$  is also a homeomorphism. Thus  $\Theta$  is a topological group isomorphism.  $\square$

*Remark 3.5.* From now on due to Theorem 3.4 we can identify  $G_{\widehat{\tau}}$  with  $G_\tau$  via the topological group isomorphism  $\Theta$  defined in (3.9). More precisely, we may identify an element  $(h, \widehat{k}) \in G_{\widehat{\tau}}$  with  $(h, k) \in G_\tau$ .

#### 4. $\tau$ -Fourier transform

In this section we study the  $\tau$ -Fourier analysis on the semi-direct product  $G_\tau$ .

We define the  $\tau$ -Fourier transform of  $f \in L^1(G_\tau)$  for a.e.  $(h, \omega) \in G_{\widehat{\tau}}$  by

$$(4.1) \quad \mathcal{F}_\tau(f)(h, \omega) := \delta(h)\mathcal{F}_K(f_h)(\omega) = \delta(h) \int_K f(h, k) \overline{\omega(k)} dk.$$

In the next theorem we prove a Parseval formula for the  $\tau$ -Fourier transform.

**Theorem 4.1.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \rtimes_\tau K$  with  $K$  abelian also let  $f \in L^1(G_\tau)$  and  $\Psi \in L^1(G_{\widehat{\tau}})$ . Define the function  $g$  for a.e.  $(h, k) \in G_\tau$  by*

$$(4.2) \quad g(h, k) := \int_{\widehat{K}} \Psi(h, \omega) \overline{\omega(k)} d\omega.$$

*Then,  $\mathcal{F}_\tau(f)_h$  belongs to  $L^\infty(\widehat{K})$  and  $g_h$  belongs to  $L^\infty(K)$  for a.e.  $h \in H$  also we have the following orthogonality relations;*

$$(4.3) \quad \int_{G_\tau} \delta(h)^{-1} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) = \int_{G_{\widehat{\tau}}} \mathcal{F}_\tau(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\widehat{\tau}}}(h, \omega),$$

$$(4.4) \quad \int_{G_\tau} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) = \int_{G_{\widehat{\tau}}} \delta(h) \mathcal{F}_\tau(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\widehat{\tau}}}(h, \omega).$$

*Proof.* Let  $f \in L^1(G_\tau)$  and  $\Psi \in L^1(G_{\widehat{\tau}})$ . It is clear that for a.e.  $h \in H$  we have  $\mathcal{F}_\tau(f)_h \in L^\infty(\widehat{K})$  and  $g_h \in L^\infty(K)$ . Using Parseval Theorem (2.6), we get

$$(4.5) \quad \int_K f(h, k) \overline{g(h, k)} dk = \int_{\widehat{K}} \mathcal{F}_K(f_h)(\omega) \overline{\Psi(h, \omega)} d\omega.$$

Thus by (4.5) we have

$$\begin{aligned}\int_{G_\tau} \delta(h)^{-1} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) &= \int_H \left( \int_K f(h, k) \overline{g(h, k)} dk \right) dh \\ &= \int_H \left( \int_{\widehat{K}} \mathcal{F}_K(f_h)(\omega) \overline{\Psi(h, \omega)} d\omega \right) dh = \int_{G_{\widehat{\tau}}} \mathcal{F}_\tau(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\widehat{\tau}}}(h, \omega).\end{aligned}$$

The same argument and also (4.5) implies (4.4).  $\square$

Due to (4.1), if  $f \in L^2(G_\tau)$  we have  $f_h \in L^2(K)$  for a.e.  $h \in H$ . Thus, according to Theorem 4.25 of [3],  $\mathcal{F}_K(f_h)$  is well-defined for a.e.  $h \in H$ . Now, in the following theorem we show that the  $\tau$ -Fourier transform (4.1) is a unitary transform from  $L^2(G_\tau)$  onto  $L^2(G_{\hat{\tau}})$ .

The next theorem can be considered as a Plancherel formula for the  $\tau$ -Fourier transform.

**Theorem 4.2.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \ltimes_\tau K$  with  $K$  abelian. The  $\tau$ -Fourier transform (4.1) on  $L^2(G_\tau)$  is an isometric transform from  $L^2(G_\tau)$  onto  $L^2(G_{\hat{\tau}})$ .*

*Proof.* Let  $f \in L^2(G_\tau)$ . Using Fubini's theorem and also Plancherel theorem (Theorem 4.25 of [3]) we have

$$\begin{aligned} \|\mathcal{F}_\tau(f)\|_{L^2(G_{\hat{\tau}})}^2 &= \int_{G_{\hat{\tau}}} |\mathcal{F}_\tau(f)(h, \omega)|^2 d\mu_{G_{\hat{\tau}}}(h, \omega) \\ &= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(f_h)(\omega)|^2 d\omega \right) \delta(h) dh \\ &= \int_H \left( \int_K |f_h(k)|^2 dk \right) \delta(h) dh \\ &= \int_H \int_K |f(h, k)|^2 \delta(h) dk dh \\ &= \int_{G_\tau} |f(h, k)|^2 d\mu_{G_\tau}(h, k) = \|f\|_{L^2(G_\tau)}^2. \end{aligned}$$

Therefore, the linear map  $f \mapsto \mathcal{F}_\tau(f)$  is an isometric in the  $L^2$ -norm. Now we show that, it is also surjective. Let  $\phi \in L^2(G_{\hat{\tau}})$ . Then, for a.e.  $h \in H$  we have  $\phi_h \in L^2(\hat{K})$ . Again using the Plancherel Theorem, there is a unique  $v^h \in L^2(K)$  such that we have  $\mathcal{F}_K(v^h) = \phi_h$ . Put  $f(h, k) := \delta(h)^{-1} v^h(k)$ , then we have  $f \in L^2(G_\tau)$ . Because,

$$\begin{aligned} \int_{G_\tau} |f(h, k)|^2 d\mu_{G_\tau}(h, k) &= \int_H \int_K |f(h, k)|^2 \delta(h) dk dh \\ &= \int_H \left( \int_K |v^h(k)|^2 dk \right) \delta(h)^{-1} dh \\ &= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(v^h)(\omega)|^2 d\omega \right) \delta(h)^{-1} dh \\ &= \int_H \left( \int_{\hat{K}} |\phi_h(\omega)|^2 d\omega \right) \delta(h) dh \\ &= \int_{G_{\hat{\tau}}} |\phi(h, \omega)|^2 d\mu_{G_{\hat{\tau}}}(h, \omega) < \infty. \end{aligned}$$

Also, for a.e.  $(h, \omega) \in G_{\hat{\tau}}$  we have

$$\begin{aligned} \mathcal{F}_\tau(f)(h, \omega) &= \delta(h) \mathcal{F}_K(f_h)(\omega) \\ &= \mathcal{F}_K(v^h)(\omega) \\ &= \phi_h(\omega) = \phi(h, \omega). \end{aligned}$$

□

Now we can prove the following Fourier inversion theorem for the  $\tau$ -Fourier transform defined in (4.1).

**Theorem 4.3.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \ltimes_\tau K$  with  $K$  abelian also let  $f \in L^1(G_\tau)$  with  $\mathcal{F}_\tau(f) \in L^1(G_{\hat{\tau}})$ . Then, for a.e.  $(h, k) \in G_\tau$  we have the following reconstruction formula;*

$$(4.6) \quad f(h, k) = \delta(h)^{-1} \int_{\hat{K}} \mathcal{F}_\tau(f)(h, \omega) \omega(k) d\omega.$$

*Proof.* Let  $f \in L^1(G_\tau)$  with  $\mathcal{F}_\tau(f) \in L^1(G_{\hat{\tau}})$ . Then, for a.e.  $h \in H$  we have  $f_h \in L^1(K)$  and  $\mathcal{F}_K(f_h) \in L^1(\hat{K})$ . Using Theorem 4.32 of [3], for a.e.  $(h, k) \in G_\tau$  we have

$$\begin{aligned} f(h, k) &= \int_{\hat{K}} \mathcal{F}_K(f_h)(\omega) \omega(k) d\omega \\ &= \delta(h)^{-1} \int_{\hat{K}} \delta(h) \mathcal{F}_K(f_h)(\omega) \omega(k) d\omega = \delta(h)^{-1} \int_{\hat{K}} \mathcal{F}_\tau(f)(h, \omega) \omega(k) d\omega. \end{aligned}$$

□

We can also define the *generalized  $\tau$ -Fourier transform* of  $f \in L^1(G_\tau)$  for a.e.  $(h, \omega) \in G_{\hat{\tau}}$  by

$$(4.7) \quad \mathcal{F}_\tau^\sharp(f)(h, \omega) := \delta(h)^{3/2} \mathcal{F}_K(f_h)(\omega_h) = \delta(h)^{3/2} \int_K f(h, k) \overline{\omega_h(k)} dk.$$

The following Parseval formula for the generalized  $\tau$ -Fourier transform can be also proved.

**Theorem 4.4.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \rtimes_\tau K$  with  $K$  abelian also let  $f \in L^1(G_\tau)$  and  $\Psi \in L^1(G_{\hat{\tau}})$ . Define the function  $g$  for a.e.  $(h, k) \in G_\tau$  by*

$$(4.8) \quad g(h, k) := \int_{\hat{K}} \Psi(h, \omega) \omega_h(k) d\omega.$$

*Then,  $\mathcal{F}_\tau^\sharp(f)_h$  belongs to  $L^\infty(\hat{K})$  and  $g_h$  belongs to  $L^\infty(K)$  for a.e.  $h \in H$  also we have the following orthogonality relations;*

$$(4.9) \quad \int_{G_\tau} \delta(h)^{-1/2} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) = \int_{G_{\hat{\tau}}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\hat{\tau}}}(h, \omega),$$

$$(4.10) \quad \int_{G_\tau} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) = \int_{G_{\hat{\tau}}} \delta(h)^{1/2} \mathcal{F}_\tau^\sharp(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\hat{\tau}}}(h, \omega).$$

*Proof.* It is easy to check that for a.e.  $h \in H$ ,  $\mathcal{F}_\tau^\sharp(f)_h$  belongs to  $L^\infty(\hat{K})$  and also  $g_h$  belongs to  $L^\infty(K)$ . Using Fubini's Theorem and also the standard Parseval formula (2.6) for a.e.  $h \in H$  we get

$$\begin{aligned} \int_K f(h, k) \overline{g(h, k)} dk &= \int_K f(h, k) \left( \int_{\hat{K}} \overline{\Psi(h, \omega)} \overline{\omega_h(k)} d\omega \right) dk \\ &= \int_{\hat{K}} \left( \int_K f_h(k) \overline{\omega_h(k)} dk \right) \overline{\Psi(h, \omega)} d\omega \\ &= \int_{\hat{K}} \widehat{f_h}(\omega_h) \overline{\Psi(h, \omega)} d\omega = \delta(h)^{-3/2} \int_{\hat{K}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \overline{\Psi(h, \omega)} d\omega. \end{aligned}$$

Now, we achieve

$$\begin{aligned} \int_{G_\tau} \delta(h)^{-1/2} f(h, k) \overline{g(h, k)} d\mu_{G_\tau}(h, k) &= \int_H \left( \int_K f(h, k) \overline{g(h, k)} dk \right) \delta(h)^{1/2} dh \\ &= \int_H \left( \int_{\hat{K}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \overline{\Psi(h, \omega)} d\omega \right) \delta(h)^{-1} dh = \int_{G_{\hat{\tau}}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \overline{\Psi(h, \omega)} d\mu_{G_{\hat{\tau}}}(h, \omega). \end{aligned}$$

The same method implies (4.10). □

If we choose  $f$  in  $L^2(G_\tau)$ , then for a.e.  $h \in H$  we have  $f_h \in L^2(K)$  and so that according to Theorem 4.25 of [3],  $\mathcal{F}_K(f_h)$  belongs to  $L^2(\hat{K})$ . In the following theorem, we show that the generalized  $\tau$ -Fourier transform (4.7) is a unitary transform from  $L^2(G_\tau)$  onto  $L^2(G_{\hat{\tau}})$ .

**Theorem 4.5.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \rtimes_\tau K$  with  $K$  abelian. The generalized  $\tau$ -Fourier transform (4.7) is an isometric transform from  $L^2(G_\tau)$  onto  $L^2(G_{\hat{\tau}})$ .*



*Proof.* Let  $f \in L^2(G_\tau)$ . Due to Proposition 3.1, Fubini's theorem and also Plancherel theorem (Theorem 4.25 of [3]) we have

$$\begin{aligned}
\|\mathcal{F}_\tau^\sharp(f)\|_{L^2(G_{\hat{\tau}})}^2 &= \int_{G_{\hat{\tau}}} |\mathcal{F}_\tau^\sharp(f)(h, \omega)|^2 d\mu_{G_{\hat{\tau}}}(h, k) \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_\tau^\sharp(f)(h, \omega)|^2 dk \right) \delta(h)^{-1} dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(f_h)(\omega_h)|^2 d\omega \right) \delta(h)^2 dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(f_h)(\omega)|^2 d\omega_{h^{-1}} \right) \delta(h)^2 dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(f_h)(\omega)|^2 d\omega \right) \delta(h) dh \\
&= \int_H \left( \int_K |f(h, k)|^2 dk \right) \delta(h) dh \\
&= \int_{G_\tau} |f(h, k)|^2 d\mu_{G_\tau}(h, k) = \|f\|_{L^2(G_\tau)}^2.
\end{aligned}$$

Now to show that the generalized  $\tau$ -Fourier transform (4.7) maps  $L^2(G_\tau)$  onto  $L^2(G_{\hat{\tau}})$ , let  $\phi \in L^2(G_{\hat{\tau}})$  be given. Then, for a.e.  $h \in H$  we have  $\phi_h \in L^2(K)$  and so that there is unique  $v^h \in L^2(K)$  with  $\mathcal{F}_K(v^h) = \phi_h$ . Put  $f(h, k) = \delta(h)^{-1/2} v^h \circ \tau_{h^{-1}}(k)$ . Then we have  $f \in L^2(G_\tau)$ , because

$$\begin{aligned}
\|f\|_{L^2(G_\tau)}^2 &= \int_{G_\tau} |f(h, k)|^2 d\mu_{G_\tau}(h, k) \\
&= \int_H \left( \int_K |f(h, k)|^2 dk \right) \delta(h) dh \\
&= \int_H \left( \int_K |v^h \circ \tau_{h^{-1}}(k)|^2 dk \right) dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(v^h \circ \tau_{h^{-1}})(\omega)|^2 d\omega \right) dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(v^h)(\omega_{h^{-1}})|^2 d\omega \right) \delta(h)^{-2} dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(v^h)(\omega)|^2 d\omega_h \right) \delta(h)^{-2} dh \\
&= \int_H \left( \int_{\hat{K}} |\mathcal{F}_K(v^h)(\omega)|^2 d\omega \right) \delta(h)^{-1} dh \\
&= \int_H \left( \int_{\hat{K}} |\phi(h, \omega)|^2 d\omega \right) \delta(h)^{-1} dh = \|\phi\|_{L^2(G_{\hat{\tau}})}^2.
\end{aligned}$$

Also, we have

$$\begin{aligned}
\mathcal{F}_\tau^\sharp(f) &= \delta(h)^{3/2} \mathcal{F}_K(f_h)(\omega_h) \\
&= \delta(h) \mathcal{F}_K(v^h \circ \tau_{h^{-1}})(\omega_h) \\
&= \mathcal{F}_K(v^h)(\omega) = \phi(h, \omega).
\end{aligned}$$

□

In the following we prove an inversion formula for the generalized  $\tau$ -Fourier transform defined in (4.7).

**Theorem 4.6.** *Let  $\tau : H \rightarrow \text{Aut}(K)$  be a continuous homomorphism and  $G_\tau = H \ltimes_\tau K$  with  $K$  abelian also let  $f \in L^1(G_\tau)$  with  $\mathcal{F}_\tau^\sharp(f) \in L^1(G_{\widehat{\tau}})$ . Then, for a.e.  $(h, k) \in G_\tau$  we have the following reconstruction formula;*

$$(4.11) \quad f(h, k) = \delta(h)^{-1/2} \int_{\widehat{K}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \omega_h(k) d\omega.$$

*Proof.* Let  $f \in L^1(G_\tau)$  and also let  $\mathcal{F}_\tau^\sharp(f) \in L^1(G_{\widehat{\tau}})$ . Due to Proposition 3.1, for a.e.  $h \in H$  we have

$$\int_{\widehat{K}} |\mathcal{F}_K(f_h)(\omega_h)| d\omega = \int_{\widehat{K}} |\mathcal{F}_K(f_h)(\omega)| d\omega_{h^{-1}} = \delta(h)^{-1} \int_{\widehat{K}} |\mathcal{F}_K(f_h)(\omega)| d\omega.$$

Thus, for a.e.  $h \in H$  we get  $\mathcal{F}_K(f_h) \in L^1(\widehat{K})$ . Now, using Theorem 4.32 of [3] and also Proposition 3.1 for a.e.  $(h, k) \in G_\tau$  we achieve

$$\begin{aligned} f(h, k) &= \int_{\widehat{K}} \mathcal{F}_K(f_h)(\omega) \omega(k) d\omega \\ &= \int_{\widehat{K}} \mathcal{F}_K(f_h)(\omega_h) \omega_h(k) d\omega_h \\ &= \delta(h) \int_{\widehat{K}} \mathcal{F}_K(f_h)(\omega_h) \omega_h(k) d\omega \\ &= \delta(h)^{-1/2} \int_{\widehat{K}} \delta(h)^{3/2} \mathcal{F}_K(f_h)(\omega_h) \omega_h(k) d\omega = \delta(h)^{-1/2} \int_{\widehat{K}} \mathcal{F}_\tau^\sharp(f)(h, \omega) \omega_h(k) d\omega. \end{aligned}$$

□

## 5. Examples

As an application we study the theory of  $\tau$ -Fourier transform for the affine group  $\mathbf{ax} + \mathbf{b}$ .

**5.1. Affine group  $\mathbf{ax} + \mathbf{b}$ .** Let  $H = \mathbb{R}_+^* = (0, +\infty)$  and  $K = \mathbb{R}$ . The affine group  $\mathbf{ax} + \mathbf{b}$  is the semi direct product  $H \ltimes_\tau K$  with respect to the homomorphism  $\tau : H \rightarrow \text{Aut}(K)$  given by  $a \mapsto \tau_a$ , where  $\tau_a(b) = ab$ . Hence the underlying manifold of the affine group is  $(0, \infty) \times \mathbb{R}$  and also the group law is

$$(5.1) \quad (a, b) \ltimes_\tau (a', b') = (aa', b + ab').$$

The continuous homomorphism  $\delta : H \rightarrow (0, \infty)$  is given by  $\delta(a) = a^{-1}$  and so that the left Haar measure is in fact  $d\mu_{G_\tau}(a, b) = a^{-2} da db$ . Due to Theorem 4.5 of [3] we can identify  $\widehat{\mathbb{R}}$  with  $\mathbb{R}$  via  $\omega(b) = \langle b, \omega \rangle = e^{2\pi i \omega b}$  for each  $\omega \in \widehat{\mathbb{R}}$  and so we can consider the continuous homomorphism  $\widehat{\tau} : H \rightarrow \text{Aut}(\widehat{K})$  given by  $a \mapsto \widehat{\tau}_a$  via

$$\begin{aligned} \langle b, \widehat{\tau}_a(\omega) \rangle &= \langle b, \omega_a \rangle \\ &= \langle \tau_{a^{-1}}(b), \omega \rangle = \langle a^{-1}b, \omega \rangle = e^{2\pi i \omega a^{-1}b}. \end{aligned}$$

Thus,  $\tau$ -dual group of the affine group again has the underlying manifold  $(0, \infty) \times \mathbb{R}$ , with  $\tau$ -dual group law given by

$$(5.2) \quad (a, \omega) \ltimes_{\widehat{\tau}} (a', \omega') = (aa', \omega + \omega'_a) = (aa', \omega + a^{-1}\omega').$$

Using Theorem 3.2, the left Haar measure  $d\mu_{G_{\widehat{\tau}}}(a, \omega)$  of  $G_{\widehat{\tau}}$  is precisely  $da d\omega$ . Now we recall that the standard dual space of the affine group which is precisely the set of all unitary irreducible representations of the affine group  $\mathbf{ax} + \mathbf{b}$ , are described via Theorem 6.42 of [3] and also Theorem 7.50 of [3] guarantee the following Plancherel formula;

$$(5.3) \quad \|\widehat{f}(\pi_+)\|_{\text{HS}}^2 + \|\widehat{f}(\pi_-)\|_{\text{HS}}^2 = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db,$$

for all measurable function  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  satisfying

$$(5.4) \quad \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db < \infty.$$

Now let  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  be a measurable function satisfying

$$(5.5) \quad \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|}{a^2} da db < \infty.$$

Then, for a.e.  $(a, \omega) \in G_{\hat{\tau}} = \mathbb{R}_+^* \ltimes_{\hat{\tau}} \mathbb{R}$  the  $\tau$ -Fourier transform of  $f$  is given by (4.1) via

$$\mathcal{F}_\tau(f)(a, \omega) = \delta(a) \hat{f}_a(\omega) = a^{-1} \int_{-\infty}^{+\infty} f(a, b) e^{-2\pi i \omega \cdot b} db,$$

and also the following Plancherel formula for the  $\tau$ -Fourier transform holds;

$$(5.6) \quad \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau(f)(a, \omega)|^2 da d\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db.$$

Therefore, we have the following proposition.

**Proposition 5.1.** *Let  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  be a measurable function satisfying (5.4). Then,*

$$(5.7) \quad \int_0^\infty \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{f(a, b) \overline{f(a, \beta)}}{a^2} e^{-2\pi i \omega (b - \beta)} db d\beta da d\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db.$$

*Proof.* Using (5.6) and also Fubini's theorem we have

$$\begin{aligned} \int_0^\infty \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{f(a, b) \overline{f(a, \beta)}}{a^2} e^{-2\pi i \omega (b - \beta)} db d\beta da d\omega &= \int_0^\infty \int_{-\infty}^{+\infty} \left| \int_{-\infty}^{+\infty} f(a, b) e^{-2\pi i \omega \cdot b} db \right|^2 \frac{da d\omega}{a^2} \\ &= \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau(f)(a, \omega)|^2 da d\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db. \end{aligned}$$

□

Due to the reconstruction formula (4.6) each measurable function  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  satisfying (5.5) with

$$(5.8) \quad \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau(f)(a, \omega)| da d\omega < \infty,$$

satisfies the following reconstruction formula;

$$\begin{aligned} f(a, b) &= \delta(a)^{-1} \int_{-\infty}^{+\infty} \hat{f}_a(\omega) e^{2\pi i \omega \cdot b} d\omega \\ &= \int_{-\infty}^{+\infty} \left( \int_{-\infty}^{+\infty} f(a, \beta) e^{-2\pi i \omega \beta} d\beta \right) e^{2\pi i \omega \cdot b} d\omega = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(a, \beta) e^{2\pi i \omega (b - \beta)} d\beta d\omega. \end{aligned}$$

For a.e.  $(a, \omega) \in G_{\hat{\tau}} = \mathbb{R}_+^* \ltimes_{\hat{\tau}} \mathbb{R}$  and also each measurable function  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  satisfying (5.4), the generalized  $\tau$ -Fourier transform of  $f$  is given by (4.7) via

$$\begin{aligned} \mathcal{F}_\tau^\#(f)(a, \omega) &= \delta(a)^{3/2} \hat{f}_a(\omega_a) \\ &= a^{-3/2} \int_{-\infty}^{+\infty} f(a, b) e^{-2\pi i \omega_a \cdot b} db = a^{-3/2} \int_{-\infty}^{+\infty} f(a, b) e^{-2\pi i \omega a^{-1} b} db. \end{aligned}$$

According to Theorem 4.5 the generalized  $\tau$ -Fourier transform satisfies the following Plancherel formula;

$$(5.9) \quad \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau^\#(f)(a, \omega)|^2 da d\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db.$$

**Proposition 5.2.** *Let  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  be a measurable function satisfying (5.4). Then,*

$$(5.10) \quad \int_0^\infty \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{f(a, b) \overline{f(a, \beta)}}{a^3} e^{-2\pi i \omega a^{-1} (b - \beta)} db d\beta da d\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a, b)|^2}{a^2} da db.$$

*Proof.* Using (5.9) and also Fubini's theorem we have

$$\begin{aligned}
\int_0^\infty \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{f(a,b)\overline{f(a,\beta)}}{a^3} e^{-2\pi i \omega a^{-1}(b-\beta)} db d\beta da d\omega &= \int_0^\infty \int_{-\infty}^{+\infty} \left| \int_{-\infty}^{+\infty} f(a,b) e^{-2\pi i \omega a^{-1}b} db \right|^2 \frac{dad\omega}{a^3} \\
&= \int_0^\infty \int_{-\infty}^{+\infty} \left| \int_{-\infty}^{+\infty} f(a,b) e^{-2\pi i \omega_a \cdot b} db \right|^2 \frac{dad\omega}{a^3} \\
&= \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau^\#(f)(a,\omega)|^2 dad\omega = \int_0^\infty \int_{-\infty}^{+\infty} \frac{|f(a,b)|^2}{a^2} dadb.
\end{aligned}$$

□

Due to the reconstruction formula (4.11) each measurable function  $f : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{C}$  satisfying (5.5) with

$$(5.11) \quad \int_0^\infty \int_{-\infty}^{+\infty} |\mathcal{F}_\tau^\#(f)(a,\omega)| dad\omega < \infty,$$

satisfies the following reconstruction formula;

$$\begin{aligned}
f(a,b) &= \delta(a)^{-1/2} \int_{-\infty}^{+\infty} \mathcal{F}_\tau^\#(f)(a,\omega) \omega_a(b) d\omega \\
&= \delta(a) \int_{-\infty}^{+\infty} \widehat{f}_a(\omega_a) \omega_a(b) d\omega \\
&= \delta(a) \int_{-\infty}^{+\infty} \left( \int_{-\infty}^{+\infty} f(a,\beta) e^{-2\pi i \omega_a \cdot \beta} d\beta \right) \omega_a(b) d\omega \\
&= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(a,\beta) e^{-2\pi i \omega_a \cdot (\beta-b)} d\beta d\omega = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(a,\beta) e^{-2\pi i \omega a^{-1}(\beta-b)} d\beta d\omega.
\end{aligned}$$

In the sequel we find the  $\tau$ -dual group of some other well-known semi-direct product groups.

**5.2. Euclidean groups.** Let  $E(n)$  be the group of rigid motions of  $\mathbb{R}^n$ , the group generated by rotations and translations. If we put  $H = \text{SO}(n)$  and also  $K = \mathbb{R}^n$ , then  $E(n)$  is the semi direct product of  $H$  and  $K$  with respect to the continuous homomorphism  $\tau : \text{SO}(n) \rightarrow \text{Aut}(\mathbb{R}^n)$  given by  $\sigma \mapsto \tau_\sigma$  via  $\tau_\sigma(\mathbf{x}) = \sigma\mathbf{x}$  for all  $\mathbf{x} \in \mathbb{R}^n$ . The group operation for  $E(n)$  is

$$(5.12) \quad (\sigma, \mathbf{x}) \ltimes_\tau (\sigma', \mathbf{x}') = (\sigma\sigma', \mathbf{x} + \tau_\sigma(\mathbf{x}')) = (\sigma\sigma', \mathbf{x} + \sigma\mathbf{x}').$$

Identifying  $\widehat{K}$  with  $\mathbb{R}$ , the continuous homomorphism  $\widehat{\tau} : \text{SO}(n) \rightarrow \text{Aut}(\mathbb{R}^n)$  is given by  $\sigma \mapsto \widehat{\tau}_\sigma$  via

$$\begin{aligned}
\langle \mathbf{x}, \mathbf{w}_\sigma \rangle &= \langle \mathbf{x}, \widehat{\tau}_\sigma(\mathbf{w}) \rangle \\
&= \langle \tau_{\sigma^{-1}}(\mathbf{x}), \mathbf{w} \rangle = \langle \sigma^{-1}\mathbf{x}, \mathbf{w} \rangle = e^{-2\pi i(\sigma^{-1}\mathbf{x}, \mathbf{w})} = e^{-2\pi i(\sigma\mathbf{x}, \mathbf{w})}.
\end{aligned}$$

where  $(\cdot, \cdot)$  stands for the standard inner product of  $\mathbb{R}^n$ . Since  $H$  is compact we have  $\delta = 1$  and so that  $d\sigma d\mathbf{x}$  is a left Haar measure for  $E(n)$ . Thus, the  $\tau$ -dual group of  $E(n)$  has underlying manifold  $\text{SO}(n) \times \mathbb{R}^n$  with the group operation

$$(5.13) \quad (\sigma, \mathbf{w}) \ltimes_{\widehat{\tau}} (\sigma', \mathbf{w}') = (\sigma\sigma', \mathbf{w} + \mathbf{w}').$$

**5.3. The Weyl-Heisenberg group.** Let  $K$  be a locally compact abelian (LCA) group with the Haar measure  $dk$  and  $\widehat{K}$  be the dual group of  $K$  with the Haar measure  $d\omega$  also  $\mathbb{T}$  be the circle group and let the continuous homomorphism  $\tau : K \rightarrow \text{Aut}(\widehat{K} \times \mathbb{T})$  via  $s \mapsto \tau_s$  be given by  $\tau_s(\omega, z) = (\omega, z \cdot \omega(s))$ . The semi-direct product  $G_\tau = K \ltimes_\tau (\widehat{K} \times \mathbb{T})$  is called the Weyl-Heisenberg group associated with  $K$  which is usually denoted by  $\mathbb{H}(K)$ . The group operation for all  $(k, \omega, z), (k', \omega', z') \in K \ltimes_\tau (\widehat{K} \times \mathbb{T})$  is

$$(5.14) \quad (k, \omega, z) \ltimes_\tau (k', \omega', z') = (k + k', \omega\omega', zz'\omega'(k)).$$

If  $dz$  is the Haar measure of the circle group, then  $dkd\omega dz$  is a Haar measure for the Weyl-Heisenberg group and also the continuous homomorphism  $\delta : K \rightarrow (0, \infty)$  given in (2.3) is the constant function 1. Thus, using Theorem 4.5 and also Proposition 4.6 of [3] and also Theorem 3.2 we can obtain the continuous homomorphism  $\widehat{\tau} : K \rightarrow \text{Aut}(K \times \mathbb{Z})$  via  $s \mapsto \widehat{\tau}_s$ , where  $\widehat{\tau}_s$  is given by  $\widehat{\tau}_s(k, n) = (k, n) \circ \tau_{s^{-1}}$  for all  $(k, n) \in K \times \mathbb{Z}$  and  $s \in K$ . Due to Theorem 4.5 of [3], for each  $(k, n) \in K \times \mathbb{Z}$  and also for all  $(\omega, z) \in \widehat{K} \times \mathbb{T}$  we have

$$\begin{aligned} \langle (\omega, z), (k, n)_s \rangle &= \langle (\omega, z), \widehat{\tau}_s(k, n) \rangle \\ &= \langle \tau_{s^{-1}}(\omega, z), (k, n) \rangle \\ &= \langle (\omega, z\overline{\omega(s)}), (k, n) \rangle \\ &= \langle \omega, k \rangle \langle z\overline{\omega(s)}, n \rangle \\ &= \omega(k) z^n \overline{\omega(s)}^n \\ &= \omega(k - ns) z^n = \langle \omega, k - ns \rangle \langle z, n \rangle = \langle (\omega, z), (k - ns, n) \rangle. \end{aligned}$$

Thus,  $(k, n)_s = (k - ns, n)$  for all  $k, s \in K$  and  $n \in \mathbb{Z}$ . Therefore,  $G_{\widehat{\tau}}$  has the underlying set  $K \times K \times \mathbb{Z}$  with the following group operation;

$$\begin{aligned} (s, k, n) \rtimes_{\widehat{\tau}} (s', k', n') &= (s + s', (k, n)\widehat{\tau}_s(k', n')) \\ &= (s + s', (k, n)(k' - n's, n')) = (s + s', k + k' - n's, n + n'). \end{aligned}$$

#### REFERENCES

1. Corwin. L., Greenleaf. F.P., *Representations of nilpotent Lie groups and their applications*, Part 1: basic theory and examples, Cambridge studies in advanced mathematics 18, Cambridge university press, 1990.
2. Dixmier. J., *C\*-Algebras*, North-Holland and Publishing company, 1977.
3. Folland. G.B., *A course in Abstract Harmonic Analysis*, CRC press, 1995.
4. Folland. G.B., *Real analysis, Modern Techniques and Their Applications*, Wiley-Inter-science publication, 1999.
5. Hewitt. E. and Ross. K. A., *Abstract Harmonic Analysis*, Springer-Verlag, Vol 1, 1963.
6. Hewitt. E. and Ross. K. A., *Abstract Harmonic Analysis*, Springer-Verlag, Vol 2, 1969.
7. Hochschild. G., *The Structure of Lie Groups*, Hpolden-day, San Francisco, (1965).
8. Kirillov. A.A., Neretin. Y. A., *Representation Theory and Non-commutative Harmonic Analysis I: Fundamental Concepts. Representations of Virasoro and Affine Algebras* v.1. (1994).
9. Lipsman. R., *Non-Abelian Fourier Analysis*, Bull. Sc. Math., 2<sup>e</sup> series, **98**, 1974, pp.209-233.
10. Murphy. G.J., *C\*-Algebras and Operator theory*, Academic Press, INC.
11. Reiter. H., and Stegeman. J. D., *Classical Harmonic Analysis*, 2nd Ed, Oxford University Press, New York, 2000.
12. Segal. I.E., *An extension of Plancherel's formula to separable unimodular groups*, Ann. of Math, **52**, 1950, pp.272-292.
13. Taylor. M.E., *Non-commutative harmonic analysis*, Mathematics surveys and Monographs, No 22, American Mathematical Society, Providence, Rhode Island, 1986.

DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, FERDOWSI UNIVERSITY OF MASHHAD (FUM), P. O. BOX 1159, MASHHAD 91775, IRAN.

E-mail address: ghaanifarashahi@hotmail.com

E-mail address: ghaanifarashahi@gmail.com